

Institute of Chemical Engineering

Adres artykułu: <https://iich.gliwice.pl/en/article/cometabolic-biodegradation-of-4-chlorophenol-by-stenotrophomonas-maltophilia-kb2-strain>

Cometabolic biodegradation of 4-chlorophenol by *stenotrophomonas maltophilia* KB2 strain

Publication date:	29.12.2016
Publication title:	Cometabolic biodegradation of 4-chlorophenol by <i>stenotrophomonas maltophilia</i> KB2 strain
Authors:	Grażyna Bartelmuś , Anna Szczotka , Hanna Kolarszyc , Agnieszka Gąsyczak , Elżbieta Szczyrba , Izabela Greń
Journal information:	Prace Naukowe Instytutu Inżynierii Chemicznej Polskiej Akademii Nauk
Tags:	phenol , cometabolism , chlorophenol

Abstract: The cometabolic biodegradation of 4-chlorophenol by *Stenotrophomonas maltophilia* KB2 strain in the presence of phenol as the sole carbon and energy source was studied. The experiments were carried out in a batch bioreactor. The tests were conducted at 30°C, pH 7 and oxygenation maintained at the level of 5–7 mg·l⁻³. The effect of changes in initial concentration of both substrates on the rate of biomass growth and substrates transformation were analyzed. The initial concentration of phenol was changed within the range of 50–300 g·m⁻³ and 4-chlorophenol within the range of 25–100 g·m⁻³. The values of growth substrate transformation yield (W) and the f coefficient, indicating the fraction of reductant from phenol oxidation used to transform 4-chlorophenol, were also estimated.

Attachments:

[Zeszyt-20-2016](#) pdf, 4.77 MB

Created at:	04.08.2025
Published by:	Artur Wojdyła
Published at:	05.08.2025 08:19

Number of downloads:	269
-----------------------------	-----

Tagi: phenol, cometabolism, chlorophenol

Metryczka

Published by:	Artur Wojdyła
Published at:	18.09.2025 13:02
Number of views:	70