

Institute of Chemical Engineering

Adres artykułu: <https://iich.gliwice.pl/en/article/kinetics-of-esterification-of-the-levulinic-acid-with-n-hexanol-n-octanol-and-2-ethylhexanol-in-the-presence-of-methanesulfonic-acid-as-a-catalyst-under-nonisothermal-conditions-1>

Kinetics of esterification of the levulinic acid with n-hexanol, n-octanol, and 2-ethylhexanol in the presence of methanesulfonic acid as a catalyst under nonisothermal conditions

Publication date:	01.01.2021
Publication title:	Kinetics of esterification of the levulinic acid with n-hexanol, n-octanol, and 2-ethylhexanol in the presence of methanesulfonic acid as a catalyst under nonisothermal conditions
Authors:	Łukasz Hamryszak , et al.
Journal information:	INTERNATIONAL JOURNAL OF CHEMICAL KINETICS

The levulinic acid was esterified with alcohol at an alcohol to acid molar ratio of 3:1, 5:1, and 10:1 in the presence of a 0.1 wt% methanesulfonic acid catalyst. During esterification, the temperature was changed linearly from 373 to 428 K and its average change was 4.5 K/min. The authors stated that reactions were of second order and that the activation energy (E) decreased from 61 to 46 kJ/mol in the following alcohol sequence: n-hexanol > n-octanol > 2-ethylhexanol. The fitting errors varied between 3.8% and 6.4%. The time of experiment carried out under nonisothermal condition is five to 15 times shorter than that conducted under isothermal conditions. A smaller number of experimental series also determines a significantly lower cost of such research. The results of such study are the precise form of the kinetic equation, which is indispensable in design and optimization of industrial-scale chemical reactors.

Metryczka

Published by:	Marek Tańczyk
Published at:	08.05.2026 14:57

Number of views:

13