

# Instytut Inżynierii Chemicznej

Adres artykułu: <https://iich.gliwice.pl/pl/artykul/transport-of-carbon-dioxide-methane-oxygen-and-nitrogen-in-a-glassy-polyimide-membrane>

## Transport of Carbon Dioxide, Methane, Oxygen and Nitrogen in a Glassy Polyimide Membrane

<b>Data publikacji:</b>	23.11.2025
<b>Tytuł publikacji:</b>	<a href="https://iich.gliwice.pl/pl/artykul/transport-of-carbon-dioxide-methane-oxygen-and-nitrogen-in-a-glassy-polyimide-membrane">Transport of Carbon Dioxide, Methane, Oxygen and Nitrogen in a Glassy Polyimide Membrane</a>
<b>Autorzy:</b>	<a href="#">Marek Tańczyk</a> , <a href="#">Aleksandra Janusz-Cygan</a> , <a href="#">Jolanta Jaschik</a> , <a href="#">Łukasz Hamryszak</a> , <a href="#">Anna Pawlaczyk-Kurek</a>
<b>Informacje o czasopiśmie:</b>	<a href="#">Molecules</a>
<b>Tagi:</b>	<a href="#">biogas</a>

Biomethane is one of the controllable Renewable Energy Sources. It may be derived from biogas, a multicomponent gas mixture, using, among others, membrane processes. The proper optimization of such a process requires the knowledge of the phenomena accompanying each specific biogas-membrane separation system. Therefore, the solubility, permeance and diffusion of CO<sub>2</sub>, CH<sub>4</sub>, O<sub>2</sub> and N<sub>2</sub> in a polyimide-based sample were described and analyzed using the Dual Mode Sorption and partial immobilization models. The parameters of the models were determined based on pure gas sorption isotherms measured gravimetrically and experimental permeances of the four gases. The membrane swelling caused by CO<sub>2</sub> was observed at temperatures of 293 and 303 K and for pressures higher than 3 bar. The adsorption of CH<sub>4</sub>, O<sub>2</sub> and N<sub>2</sub> in the fractional free volume (FFV) has a dominant (>50%) share in their total solubility in the entire pressure range. This makes them sensitive to the presence of CO<sub>2</sub>, whose affinity is the strongest towards the tested polyimide-based sample. The diffusion of O<sub>2</sub> is the fastest which makes it competitive with CO<sub>2</sub> in permeation through the membrane, despite its low solubility. The ideal CO<sub>2</sub>/O<sub>2</sub> selectivity is thus relatively low (2.3–5.1). Methane, which is competitive in solubility compared to CO<sub>2</sub>, was found to diffuse the slowest and as a result, it is also the slowest permeating gas. This translates into the very high CO<sub>2</sub>/CH<sub>4</sub> ideal selectivity (33–95.7), which is, however, strongly dependent on temperature and pressure.

Tagi: biogas

# Metryczka

<b>Data opublikowania:</b>	25.11.2025 10:23
<b>Ostatnio zaktualizował:</b>	Marek Tańczyk
<b>Data ostatniej aktualizacji:</b>	26.01.2026 14:49
<b>Liczba wyświetleń:</b>	231