

# Instytut Inżynierii Chemicznej

Adres artykułu: <https://iich.gliwice.pl/pl/artykul/lipase-immobilized-on-mcfs-as-biocatalysts-for-kinetic-and-dynamic-kinetic-resolution-of-sec-alcohols>

## Lipase immobilized on MCFs as biocatalysts for kinetic and dynamic kinetic resolution of sec-alcohols

<b>Data publikacji:</b>	20.04.2021
<b>Tytuł publikacji:</b>	<a href="https://iich.gliwice.pl/pl/artykul/lipase-immobilized-on-mcfs-as-biocatalysts-for-kinetic-and-dynamic-kinetic-resolution-of-sec-alcohols">Lipase immobilized on MCFs as biocatalysts for kinetic and dynamic kinetic resolution of sec-alcohols</a>
<b>Autorzy:</b>	<a href="#">Katarzyna Maresz</a> , <a href="#">Andrzej B. Jarzębski</a>
<b>Informacje o czasopiśmie:</b>	<a href="#">Catalysts</a>

Dynamic kinetic resolution (DKR) is one of the most attractive methods for enantioselective synthesis. In the reported studies, lipase B from *Candida antarctica* (CALB) immobilized on siliceous mesoporous cellular foams (MCF) functionalized with different hydrophobic groups, and two ruthenium complexes with substituted cyclopentadienyl ligands were investigated as catalysts for the chemoenzymatic DKR of (rac)-1-phenylethanol, using Novozym 435 as a benchmark biocatalyst. Studies on the (rac)-1-phenylethanol transesterification reaction showed that CALB supported on MCFs grafted with methyl groups is a promising biocatalyst and isopropenyl acetate is a preferable acylation agent. Both Ru-complexes activated by K<sub>3</sub>PO<sub>4</sub> or t-BuOK, proved to be effective catalysts of the racemization reaction. The final DKR experiments using all catalysts combinations singled out, gave 96% conversion, and (R)-1-phenylethyl acetate enantiomeric excess of 98% in 8 h using K<sub>3</sub>PO<sub>4</sub> activator.

## Metryczka

<b>Opublikował w BIP:</b>	Marek Tańczyk
<b>Data opublikowania:</b>	08.05.2026 15:06
<b>Liczba wyświetleń:</b>	19