

# Instytut Inżynierii Chemicznej

Adres artykułu: <https://iich.gliwice.pl/pl/artykul/effect-of-nickel-as-stress-factor-on-phenol-biodegradation-by-stenotrophomonas-maltophilia-kb2>

## Effect of Nickel as Stress Factor on Phenol Biodegradation by *Stenotrophomonas maltophilia* KB2

<b>Data publikacji:</b>	14.10.2021
<b>Tytuł publikacji:</b>	<a href="https://iich.gliwice.pl/pl/artykul/effect-of-nickel-as-stress-factor-on-phenol-biodegradation-by-stenotrophomonas-maltophilia-kb2">Effect of Nickel as Stress Factor on Phenol Biodegradation by <i>Stenotrophomonas maltophilia</i> KB2</a>
<b>Autorzy:</b>	<a href="#">Anna Szczotka</a> , <a href="#">Agnieszka Gąszczak</a> , <a href="#">Elżbieta Szczyrba</a> , <a href="#">et al.</a>
<b>Informacje o czasopiśmie:</b>	<a href="#">Materials</a>

This study focuses on the phenol biodegradation kinetics by *Stenotrophomonas maltophilia* KB2 in a nickel-contaminated medium. Initial tests proved that a nickel concentration of 33.3 mg·L<sup>-1</sup> caused a cessation of bacterial growth. The experiments were conducted in a batch bioreactor in several series: without nickel, at constant nickel concentration and at varying metal concentrations (1.67–13.33 g·m<sup>-3</sup>). For a constant Ni<sup>2+</sup> concentration (1.67 or 3.33 g·m<sup>-3</sup>), a comparable bacterial growth rate was obtained regardless of the initial phenol concentration (50–300 g·m<sup>-3</sup>). The dependence  $\mu = f(S_0)$  at constant Ni<sup>2+</sup> concentration was very well described by the Monod equations. The created varying nickel concentrations experimental database was used to estimate the parameters of selected mathematical models, and the analysis included different methods of determining metal inhibition constant K<sub>IM</sub>. Each model showed a very good fit with the experimental data (R<sup>2</sup> values were higher than 0.9). The best agreement (R<sup>2</sup> = 0.995) was achieved using a modified Andrews equation, which considers the metal influence and substrate inhibition. Therefore, kinetic equation parameters were estimated:  $\mu_{max} = 1.584 \text{ h}^{-1}$ ,  $K_S = 185.367 \text{ g}\cdot\text{m}^{-3}$ ,  $K_{IS} = 106.137 \text{ g}\cdot\text{m}^{-3}$ ,  $K_{IM} = 1.249 \text{ g}\cdot\text{m}^{-3}$  and  $n = 1.0706$

## Metryczka

<b>Opublikował w BIP:</b>	Marek Tańczyk
---------------------------	---------------

<b>Data opublikowania:</b>	08.05.2026 15:21
<b>Liczba wyświetleń:</b>	6